

Solid-solid volume collapse transitions are zeroth order

S. Bustingorry,¹ E. A. Jagla,¹ and J. Lorenzana²

¹*Consejo Nacional de Investigaciones Científicas y Técnicas,
Centro Atómico Bariloche, 8400 Bariloche, Argentina*

²*Center for Statistical Mechanics and Complexity, INFM, Dipartimento di Fisica,
Università di Roma La Sapienza, P. Aldo Moro 2, 00185 Roma, Italy*

(Dated: February 2, 2008)

We present an exactly solvable non-linear elastic model of a volume collapse transition in an isotropic solid. Integrity of the lattice through the transition leads to an infinite-range density-density interaction, which drives classical critical behavior. Nucleation is forbidden within a pressure window leading to intrinsic hysteresis and an unavoidable discontinuity of the thermodynamic potential (zeroth order transition). The window shrinks with increasing temperature ending at a critical point at a temperature related to the shear modulus. Mixed phases behave non-extensively and show negative compressibility. We discuss the implications for Ce, SmS, and related systems.

PACS numbers: 64.70.Kb, 68.35.Rh, 71.28.+d, 71.10.Hf

Room temperature γ Cerium shows a well known transition to isostructural α Cerium at ~ 0.8 GPa with a 17% reduction of the volume. Similar (though not necessarily isostructural) transitions occur in other f -electron systems[1, 2]. Volume-collapse transitions were also predicted to occur in colloidal systems[3] and close to first order electronic transitions when mesoscopic inhomogeneous electronic states are frustrated by the long-range Coulomb interaction[4].

Consider one of these systems which is driven across the volume-collapse transition by adiabatically changing the average volume. In analogy with the van der Waals liquid-gas transition one expects a range of average volume values for which there is phase coexistence at a constant equilibrium pressure. This presents an interesting nucleation problem. Within purely energetic considerations the mixed phase consists of a region of the sample with the small unit cell volume and a region with the large unit cell volume separated by a flat surface so as to minimize the interface boundary energy. The lattice mismatch at the interface can be seen as a line of dislocations. However dislocations are very difficult to nucleate in a solid since they require the breaking of chemical bonds. Indeed it is customary to assume on solid-solid transitions that the lattice integrity is preserved[5, 6, 7]. Under this hypothesis the thermodynamic ground state is frustrated from the onset, and the question arises of how the system minimizes its free energy, preserving the integrity constraints. To answer this question we solve a non-linear elastic model of a volume-collapse transition in an isotropic solid assuming lattice integrity (no dislocations). The latter condition leads to the well known St. Venant compatibility constraints[8] among the different components of the strain tensor and allows the free energy to be expressed as a function of the order parameter (OP) alone, with long-range interactions[5, 6, 7]. This kind of approach has been shown to lead to anomalous nucleation in volume conserving transitions[7].

We find that compatibility constraints induce a distance-independent density-density interaction and thus mean-field (MF) theory becomes exact and the model results to be trivially solvable at all temperatures in the absence of short-range interactions. Critical exponents are MF like even in the presence of short-range interactions. In the mixed phase the system behaves non extensively and violates usual stability criteria for extensive systems. In particular, it possesses negative compressibility. For an experiment in which the pressure acts as the control variable we show that usual nucleation is forbidden within a finite pressure window. This pressure window implies an intrinsic hysteresis loop in a pressure controlled experiment. Outside the pressure window the transition can occur, but only irreversibly with an intrinsic jump of the thermodynamic potential. Since the discontinuity occurs in the thermodynamic potential itself rather than in its first derivative the transition can be called of zeroth order (instead of first) as done with similar phenomena in gravitational systems[9]. The extent of the hysteresis loop reduces as a function of temperature, vanishing at a critical point. The temperature of the critical point is related to the shear modulus of the material.

Model The OP is taken to be the dilation strain $e_1 \equiv \sum_{i=1}^d \epsilon_{ii}/\sqrt{d}$ where ϵ_{ij} are the components of the infinitesimal strain tensor[8] ϵ and d is the dimensionality of the system. Our starting point is a coarse grained[10] elastic Hamiltonian $H \equiv H_1 + H_h$ with H_1 a non-linear Ginzburg-Landau like double-well term:

$$H_1 = \int d^d x \left(a e_1 - \frac{b}{2} e_1^2 + \frac{c}{4} e_1^4 + \frac{\kappa}{2} |\nabla e_1|^2 \right), \quad (1)$$

The remaining $d(d+1)/2 - 1$ independent components of the strain tensor give the harmonic contribution:

$$H_h = \mu \int d^d x \sum_{i,j=1}^d \tilde{\epsilon}_{ij}^2. \quad (2)$$




FIG. 1: (a) $\bar{\phi}$ -dependent common tangent construction. The cases $\bar{\phi} = -0.5$ and 0.2 are illustrated ($m = 1$). The triangles indicate the minimum value of h attainable for the corresponding values of $\bar{\phi}$. (b) Analytical results for $\bar{h}(\bar{\phi})$ as a function of $\bar{\phi}$ for $\gamma = 0$ and different values of m , as indicated. For $m > 2$ the solution is $\bar{h}(\bar{\phi}) = h_0(\bar{\phi})$.

where μ is the shear modulus of the material, and the (traceless) ‘accommodation strains’ tensor $\tilde{\epsilon}_{ij} \equiv \epsilon_{ij} - \delta_{ij}e_1/\sqrt{d}$ has been defined.

Zero temperature theory In the absence of dislocations and in order to be physical the strain tensor must fulfill the St. Venant compatibility condition[8]:

$$\text{curl curl } \epsilon = \mathbf{0} \quad (3)$$

which leads to constraints between e_1 and $\tilde{\epsilon}_{ij}$. A uniform strain satisfies Eq. (3) trivially and therefore it is unconstrained. Thus the $\mathbf{k} = \mathbf{0}$ term in the Fourier decomposition of H_h , corresponding to *uniform* accommodation strains, can be minimized by choosing $\tilde{\epsilon}_{ij}(\mathbf{k} = \mathbf{0}) = 0$. For $\mathbf{k} \neq \mathbf{0}$ the accommodation strains can be eliminated in favor of e_1 by minimizing H_h with respect to $\tilde{\epsilon}_{ij}$, and enforcing the constraints in Eq. (3) by Lagrange multipliers[6]. The resulting equilibrium equations together with Eq. (3) allow Eq. (2) to be written as a function of e_1 alone:

$$H_h = (d-1)\mu \frac{1}{L^d} \sum_{\mathbf{k} \neq \mathbf{0}} e_1(\mathbf{k})e_1(-\mathbf{k}).$$

Here L is the linear dimension of the system. A crucial point is that the $\mathbf{k} = \mathbf{0}$ term is excluded from the sum. We can extend the sum to all $\mathbf{k}$ by explicitly subtracting the missing term. We obtain in real space:

$$\begin{aligned} H_h/[(d-1)\mu] &= \int d^d x [e_1(\mathbf{x}) - \bar{e}_1]^2 \\ &= \int d^d x \int d^d x' \left[\delta(\mathbf{x} - \mathbf{x}') - \frac{1}{L^d} \right] e_1(\mathbf{x})e_1(\mathbf{x}') \end{aligned} \quad (4)$$

with $\bar{e}_1 \equiv \int d^d \mathbf{x} e_1/L^d$. The accommodation strains induce a local term that increases the stiffness at all wave vectors, and a distance independent ‘ferro’ like interaction in e_1 that decreases the uniform stiffness alone, i.e., the stiffness at finite wave vector is larger than at zero wave vector, a well known result that holds also for linear elastic solids[10]. A similar result has been obtained in $d = 2$ in the context of the dislocation mediated melting problem close to an isostructural critical point[11].

We define $\phi \equiv e_1 \sqrt{c/b}$, $m \equiv 4(d-1)\mu/b$ and $\gamma \equiv 2\kappa/b$. Up to an additive constant the coarse grained energy density h [from Eqs. (1) and (4)] in units of the height of the double well barrier [$b^2/(4c)$] reads:

$$h(\phi, \bar{\phi}) = h_0(\phi) + \gamma|\nabla\phi|^2 + m(\phi - \bar{\phi})^2 \quad (5)$$

with $h_0(\phi) \equiv (1 - \phi^2)^2$ being the double well in the dimensionless volume ϕ . The term linear in the OP has been eliminated by redefining the origin of pressure. The parameter γ fixes the width of the interface between the low and high volume phases in a mixed state.

We first discuss the $\gamma = 0$ case and generalize for a finite γ below. The energy density Eq. (5) couples the values of ϕ at different spatial positions only through $\bar{\phi}$, namely it corresponds effectively to a MF. We stress however, that this is the *exact* energy of the model.

Mixed phase solutions for a constrained total volume obey the familiar common tangent construction: $\partial h(\phi, \bar{\phi})/\partial \phi|_{\phi=\phi_-} = \partial h(\phi, \bar{\phi})/\partial \phi|_{\phi=\phi_+} = [h(\phi_-, \bar{\phi}) - h(\phi_+, \bar{\phi})]/(\phi_- - \phi_+)$ where $\phi_{\pm} = \pm\sqrt{(1-m/2)}$ and the $+$ ($-$) corresponds to the OP of the expanded (collapsed) phase. The only difference with a conventional phase separation computation is that the construction is $\bar{\phi}$ -dependent because h depends on the global variable $\bar{\phi}$ [see Fig. 1(a)]. This dependence makes the total energy of the mixture non additive. In Fig. 1(b) we report the resulting average energy density $\bar{h}(\bar{\phi})$. For $m > 2$ the mixed solution does not exist and the system remains always uniform. We have checked these results with direct analytical computations assuming specific shapes for the coexisting phases and different boundary conditions. In all cases the maximum shear strain is of the same order as the relative volume mismatch. As long as the volume mismatch is small (a few percent) the shear strain will behave elastically. For large volume mismatch glide of atomic planes may take place and our theory becomes approximate. In any case the transition will be irreversible.

For a pressure controlled experiment the physical pressure is given by $P = p\sqrt{b^3/(16cd)} - a/\sqrt{d}$ with p the dimensionless pressure: $p = -\partial \bar{h}(\bar{\phi})/\partial \bar{\phi}$ at $T = 0$. In the mixed phase the free energy has negative curvature so the compressibility is negative [Fig. 1(b)]. This thermodynamic state is stable at constant volume with the pinned boundary conditions of Ref. [12]. This violation of usual stability criteria is common in non-extensive systems[9].

In Fig. 2 we show the values of $\bar{\phi}$ and the Gibbs free energy density $g(p) \equiv \bar{h}[\bar{\phi}(p)] + p\bar{\phi}(p)$ as a function of p

for two values of m . Controlling the total volume the system would follow the continuous line. The straight portion of this solution in Fig. 2(a) corresponds to the mixed phase. Consider now a pressure controlled experiment for $m < 4/3$ [Fig. 2(a) and (c)]. Starting from the high volume phase at low pressure and increasing p the system remains uniform until $p^\uparrow$, given by the point in which the mixed solution line intersects the uniform solution. At this point the small volume phase nucleates spontaneously since $\gamma = 0$. It is easy to show that the mixed state is mechanically unstable at constant pressure due to the negative compressibility, so the volume collapses and the system jumps to the lower branch of the curve following the vertical line. Decreasing the pressure the process is reverted and the volume suddenly expands at $p^\downarrow (= -p^\uparrow)$ producing the hysteresis loop indicated in the figure. A conventional first-order phase transition driven by pressure at a very low rate (so that nucleation occurs) would follow a vertical line at $p = 0$ in the top panel. This is not possible here because the infinite range interaction makes the critical drop radius[10] diverge, and the nucleation energy-barrier to scale as the volume of the system.

In the case $m > 4/3$ the compressibility of the uniform solution diverges at the spinodal points [full dots in Fig. 2 (b) and (d)]. The transition proceeds by a $\mathbf{k} = \mathbf{0}$ instability at $p^\uparrow = -p^\downarrow = 8/(3\sqrt{3})$, independent of m and with no energy barrier. The lower panels in Fig. 2 show that the transition involves an unavoidable jump in the thermodynamic potential for all $m \neq 0$ justifying calling this transition ‘zeroth order’[9].

Finite temperature theory Using MF theory it is straightforward to obtain the exact partition function for $\gamma = 0$, and the exact $\bar{\phi} - p$ dependence at different temperatures. In Fig. 3 we show the isotherms in the $\bar{\phi} - p$ plane for $m = 0.6$ for different values of a dimensionless




FIG. 2: Mean volume $\bar{\phi}$ and Gibbs free energy density for $m = 0.6$ (left) and $m > 2$ (right). Dotted lines are the uniform solution. Continuous lines are the solutions that minimize Eq. (5) for each value of $\bar{\phi}$. The arrows and dashed lines indicate the hysteresis cycle in pressure controlled experiments. Full dots correspond to $p^\uparrow, p^\downarrow$.

temperature t (units are restored below). The pressure width (Δp) and volume width ($\Delta \bar{\phi}$) of the hysteresis loop is reduced at finite temperatures up to the point where it vanishes, which determines the critical point coordinates ($p_c = 0, t_c(m)$). The pressure width as a function of temperature determines a wedge of forbidden nucleation [shown in Fig. 3(b) for different values of m] around the critical pressure $p_c = 0$.

For $m < 4/3$ the singularities at $p = \pm p^\uparrow$ of the $T = 0$ isotherm become rounded at finite temperature [Fig. 3(a)] due to the thermally induced spatial fluctuations of the OP. This implies that the compressibility will show precursor effects for all $m > 0$, diverging at the transition.

The solution in the case of finite γ and m is given implicitly by $\bar{\phi} = \bar{\phi}_0(p + m\bar{\phi})$ where $\bar{\phi}_0(p)$ is the equation of state of the same model except that $m = 0$, i.e., the well studied ϕ^4 -model which has the universality class of the short-range Ising model[10]. If $m > 4/3$ the qualitative description of the forbidden nucleation region are completely equivalent to the $\gamma = 0$ case. For $m < 4/3$ a region of normal nucleation and growth appears around the wedge of forbidden nucleation at low temperatures and the divergence of the compressibility gets a cutoff. These effects never reach the critical point. Both for $\gamma = 0$ and $\gamma > 0$ we find that critical exponents are given exactly by MF. This is also in agreement with the $d = 2$ theoretical analysis of Ref. [11].

Comparison with experiments A wedge where nucleation does not occur is observed in Ce[1, 2], SmS[13] and related systems. Interestingly it is also observed in a volume-collapse transition in amorphous ice[14]. However without detailed studies it is difficult to be sure that it does not arise from usual hysteresis due to a finite driving rate, as occurring in other first order transitions. Other indicators however, point more strongly to the applicability of our theory.

In agreement with the present theory MF critical exponents have been observed[2]. The connection between MF critical behavior and long-range strains has long been




FIG. 3: (a) $\bar{\phi}$ vs. p for $m = 0.6$ and different values of the dimensionless temperature t , as indicated. (b) $t - p$ phase diagram for $\gamma = 0$ and various m values (indicated in the figure). The interior of the wedge (for each value of m) is a region in which nucleation is forbidden.

suspected[2].

In contrast with usual first-order phase transitions, where hysteresis is not intrinsic, one can define here a critical exponent for the pressure width of the hysteresis loop: $\Delta p \propto |t - t_c|^{3/2}$. We are not aware of detailed measures of this critical exponent, however the curvature of the boundaries found in Ref. [13] for Gd doped SmS are compatible with this exponent. We hope our work will stimulate further experimental work.

Also in contrast with usual first-order transitions, critical behavior (although classic like) can be easily observed far from the critical point. The divergence of the compressibility discussed above implies that the bulk modulus vanishes as $|p - p^\dagger|^{1/2}$ and jumps discontinuously to a finite value for $p > p^\dagger$. Decreasing pressure the jump and the square root singularity change side and occur at $p^\dagger$. Similar behavior is found as a function of temperature.

An anomaly measured in YbInCu₄ on cooling resembles our prediction when plotted in an appropriate scale (see inset of Fig. 4 in Ref. [15]). This also explains the appearance of precursor effects which have been very puzzling in the mixed valence literature and have led some authors to interpret the transition as second order[15].

In mixed valence systems the double well is attributed to the anomalous contribution of f -electrons to the crystal binding[16]. For a realistic description it may be necessary to take the parameters in Eq. (1) to depend on temperature to take into account temperature effects of the f -electrons. In fact a trivial linear and quadratic temperature dependence of a fits our symmetric phase diagram to the asymmetric forms of Ce[1, 2] and SmS[2, 13].

The temperature dependence of a does not affect the critical temperature, but b does. Indeed in current explanations of the critical point in f -electron systems this electronic effect is crucial whereas the shear rigidity does not play a relevant role[16]. To check the relevance of the temperature scale set by the shear modulus on determining real values of T_c we take the opposite point of view, i.e., we neglect the temperature dependence of the coefficients in Eq. (1) and take into account the effect of the shear rigidity alone. Restoring dimensions, the critical temperature is given by: $k_B T_c = t_c(m)v_0 b^2/(4c)$ where v_0 is the minimum volume in the system allowed to fluctuate independently. For the case $\gamma = 0$ this volume should be assimilated to the atomic volume. In the case $\gamma > 0$, this volume can be greater depending on the strength of the short distance interactions parametrized by γ . In any case we expect it to be of the order of a few atomic volumes. The parameters b , c and m can be fixed by fitting the observed pressure and volume with of the hysteresis loop at a given temperature and from the measured value of the shear modulus[1, 17]. Within the experimental uncertainties this procedure gives the right value of T_c for Ce ($\sim 550\text{K}$ [1, 2]) and SmS ($\sim 750\text{K}$ [2, 13]) if we assume $v_0 \sim 2v_f$ where v_f is the volume associated with one formula unit. Since

the relation between v_0 and v_f go beyond our model we cannot exclude temperature effects on the coefficients of Eq. (1) contributing to T_c [16] however the fact that we obtain the right order of magnitude with a v_0 on the order of v_f implies that the scale set by the shear modulus can not be neglected in the determination of T_c .

In conclusion we have presented an exactly solvable model of an isostructural volume collapse transition, the solid-state analog of the van der Waals transition. Unlike the fluid counterpart the thermodynamics is very anomalous due to the presence of infinite range interactions. In particular, the transition is intrinsically irreversible with a discontinuity of the thermodynamic potential making of it a laboratory example of a class of phase transitions, namely zero order transitions, first discussed in the context of gravitational systems[9].

Despite the simplicity of our model, which assumes an isotropic solid, we believe the present features are quite robust. Indeed our findings are in good agreement with the behavior of real anisotropic systems[1, 2, 13, 15]. In these mixed valence systems the unexpected thermodynamics is a consequence of the interplay between strong correlations, determining the double well potential[16], and long-range strains. The present approach can be seen as a first step to understand this interplay in more complex non-isostructural transitions like the magnetoresistant manganites[4] which also show volume mismatch among different phases and irreversibility[18].

-
- [1] *Handbook on the physics and chemistry of Rare Earths*, edited by K. A. Gschneidner, Jr. and L. Eyring (North-Holland, Amsterdam, 1979), Vol. I.
 - [2] J. M. Lawrence, P. S. Riseborough, and R. D. Parks, Rep. Prog. Phys. **44**, 1 (1981).
 - [3] P. Bolhuis and D. Frenkel, Phys. Rev. Lett. **72**, 2211 (1994).
 - [4] J. Lorenzana, C. Castellani, and C. Di Castro, Phys. Rev. B **64**, 235127 (2001); **64**, 235128 (2001); Europhys. Lett. **57**, 704 (2002).
 - [5] G. R. Barsch and J. A. Krumhansl, Phys. Rev. Lett. **53**, 1069 (1984).
 - [6] S. Kartha *et al.*, Phys. Rev. B **52**, 803 (1995). K. Ø. Rasmussen *et al.*, Phys. Rev. Lett. **87**, 055704 (2001).
 - [7] W. Klein *et al.*, Phys. Rev. Lett. **88**, 085701 (2002).
 - [8] D. Chandrasekharaiah and L. Debnath, *Continuum Mechanics* (Academic Press, San Diego, 1994).
 - [9] H. J. de Vega and N. Sancés, Nuc. Phys. **625**, 409 (2002) and references therein.
 - [10] P. M. Chaikin and T. C. Lubensky, *Principles of Condensed Matter Physics* (University Press, Cambridge, 1995).
 - [11] T. Chou and D. R. Nelson, Phys. Rep. **53**, 2560 (1996).
 - [12] D. J. Bergman and B. I. Halperin, Phys. Rev. B **13**, 2145 (1976).
 - [13] V. I. Rashchupkin *et al.*, Sov. Phys. Solid State **20**, 463 (1978).

- [14] O. Mishima, J. Chem. Phys. **100**, 5910 (1994).
- [15] B. Kindler *et al.*, Phys. Rev. B **50**, 704 (1994).
- [16] J. W. Allen and R. M. Martin, Phys. Rev. Lett. **49**, 1106 (1982). B. Johansson *et al.*, *ibid.* **74**, 2335 (1995).
- [17] Tu Hailing, G. A. Saunders and H. Bach, Phys. Rev. B **29**, 1848 (1984).
- [18] D. E. Cox *et al.*, Phys. Rev. B **57**, 3305 (1998).